|                               | •                             |                    |                                |
|-------------------------------|-------------------------------|--------------------|--------------------------------|
| Treatment                     | Days required for budding (d) | Induction rate (%) | Average number of buds (block) |
| MS                            | -                             | 0                  | 0                              |
| $B_1N_1$                      | 30-35                         | 20.0 e             | 9 e                            |
| $B_1N_2$                      | 29-34                         | 24.4 d             | 11 d                           |
| B <sub>1</sub> N <sub>3</sub> | 28-33                         | 22.2 de            | 10 de                          |
| $B_2N_1$                      | 25-32                         | 28.9 cd            | 13 cd                          |
| $B_2N_2$                      | 23-28                         | 42.2 b             | 19 b                           |
| $B_2N_3$                      | 22-28                         | 31.1 c             | 14 c                           |
| $B_3N_1$                      | 20-24                         | 24.4 d             | 11 d                           |
| $B_3N_2$                      | 20-25                         | 33.3 c             | 15 c                           |
| B <sub>3</sub> N <sub>3</sub> | 22-26                         | 51.1 a             | 31 a                           |

| Supplementary Table 1c | Effect of Different Hormone Ratio on Induction of Pineapple |
|------------------------|---|
| Supplementary rable is | Effect of Different Hornone Ratio on induction of Pineappie |

Induction rates are average values. Different letters indicate significant differences (p < 0.05).